

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> :  C07K 3/04	A1	(11) International Publication Number: WO 93/20098 (43) International Publication Date: 14 October 1993 (14.10.93)
<p>(21) International Application Number: PCT/US93/02846</p> <p>(22) International Filing Date: 26 March 1993 (26.03.93)</p> <p>(30) Priority data: 07/865,368 7 April 1992 (07.04.92) US</p> <p>(71) Applicant: THE SCRIPPS RESEARCH INSTITUTE [US/US]; 10666 North Torrey Pines Road, La Jolla, CA 92037 (US).</p> <p>(72) Inventors: KENT, Stephen, B., H. ; 2766 Costebelle Drive, La Jolla, Ca 92037 (US). SCHNOLZER-RACKWITZ, Martina ; 8533-C Villa La Jolla Drive, La Jolla, CA 92037 (US).</p>		<p>(74) Agents: FITTING, Thomas et al.; The Scripps Research Institute, 10666 North Torrey Pines Road, TPC-8, La Jolla, CA 92037 (US).</p> <p>(81) Designated States: AU, CA, JP, KR, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report.</i></p>
<p>(54) Title: PROCESS FOR PREPARING MODIFIED PROTEINS</p> <p>(57) Abstract</p> <p>Process for the preparation of modified proteins comprising the coupling of a first peptide segment having a haloacyl group at the N-terminus thereof with a second peptide segment having a carbonylthiol group at the C-terminus thereof are disclosed. Novel modified proteins produced by the process are also disclosed.</p> <p>Serial No.: 09/144,838 Inventor: Siani et al. Attny Docket No. GRFN-020/01US</p> <p style="text-align: right;">F3</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

-1-

PROCESS FOR PREPARING MODIFIED PROTEINSBACKGROUND OF THE INVENTION5     FIELD OF THE INVENTION

This invention is in the area of modified biomolecules and methods of making such modified biomolecules. More particularly, this invention relates to protein  
10     engineering by chemical means to produce modified proteins where one or more peptide bonds are substituted by non-peptide linkage and one or more encoded amino acids may be replaced by unnatural amino acids or amino acid analogs or any other non-coded  
15     structure.

RELATED ART

Numerous attempts have been made to develop a  
20     successful methodology for synthesizing modified biomolecules such as proteins, glycoproteins, nucleotides, polysaccharides, and other biopolymers. Such modified biomolecules are invaluable for study of structure-activity relationships of native  
25     biomolecules and there is a growing number of commercial applications of these molecules for diagnostic or therapeutic purposes.

Structural modification of proteins and peptides,  
30     normally referred to as "protein engineering" involves the rationally designed alteration of structure with the aim of understanding protein structure and function and of creating a protein with new desirable properties. In the past, this has been principally  
35     carried out by site-directed mutagenesis or other

- 2 -

techniques involving genetic manipulation. The major drawbacks of these prior art approaches are that amino acids replacing native amino acids are those that must be coded genetically. As a result, other structural variants such as unnatural amino acids or amino acid analogs cannot be introduced in the protein backbone. However, recent findings (Ellman, et al., *Science*, 255:197, 1992; Noren, et al., *Science*, 24:182, 1989) would allow unnatural amino acids or amino acid analogs to be incorporated into proteins in a site-specific manner. In this approach, a codon encoding an amino acid to be replaced is substituted by the nonsense codon TAG by means of oligonucleotide-directed mutagenesis. A suppressor tRNA directed against this codon is then chemically aminoacylated with the desired unnatural amino acid. Addition of the amino acylated tRNA to an *in vitro* protein synthesizing system programmed with the mutagenized DNA directs the insertion of the prescribed amino acid into the protein at the target site. Taking the enzyme T4 lysozyme, the above authors, incorporated a wide variety of amino acid analogs into the enzyme at alanine 82 position with a few exceptions, for example, of D-alanine not being incorporated.

While Schultz's approach partially solves problems posed by biosynthetic protein engineering, it does not allow the alteration of the protein backbone at more than one target site to incorporate two or more different non-coded structural units. Also, by the very nature of the system, i.e., the fact that it relies upon a living system to produce the engineered protein, many substitutions or alterations, such as those which would result in a lethal mutation, cannot

- 3 -

be done. The chemical synthesis would overcome the shortcomings left by the Schultz techniques. (reviewed by R. E. Offord, *Protein Eng.*, 1:151, 1987). However, chemical synthesis is fraught with many difficulties such as the need of protection of unwanted reactive groups.

Overall, there is a definite need for a simple and efficient method for making a modified protein which possesses desired properties. The present invention addresses such need and provides novel modified proteins.

#### SUMMARY OF THE INVENTION

This invention provides new and useful modified biomolecules. It also provides a new process for producing such modified biomolecules. In general, the modified biomolecules of this invention comprise two molecular segments, each selected from peptides, pseudopeptides, or non-peptide linear molecules linked through a non-amido linkage to form a peptide or pseudopeptide backbone, wherein one or the segment contains at least one non-coded structural unit and the non-coded structural unit does not form a part of the non-amido linkage. The chemical bonding of the two segments is by means of terminal reactive groups on one segment which react with reactive groups of the other segment molecule.

The process of this invention provides a directed ligation of the two molecular segments to create a desired bond at the ligation point(s) and comprises the steps of:

- 4 -

- 5           a.    providing a first segment having at least one non-coded structural unit and attaching a first chemoselective synthon to the first segment at the terminal position thereof;
- 10           b.    providing a second segment optionally containing at least one non-coded structural unit, and second chemoselective synthon at the terminal position thereof, the second chemoselective synthon being complementary to the first chemoselective synthon of the first segment; and
- 15           c.    ligating the first segment and the second segment, whereby the first synthon of the first segment and the second synthon of the second segment forms a non-peptide linkage, wherein the first segment and the second segment are each selected from peptides, pseudopeptides, or non-peptide linear molecules, provided that both segments are not non-peptide linear molecules at the same time.
- 20
- 25           The above sequence a-c can be repeated by using a first modified biomolecule as the first segment to which a second segment or a second biomolecule is ligated. The present process also may include the step of ligating additional segments with the first and second segments which have been provided with additional terminal synthons that are compatible with the first and second synthons and chemoselective to synthons of the additional segments.
- 30

- 5 -

The present invention is therefore applicable in the chemical synthesis of various protein conjugates, such as proteins with reporter molecules, radionuclides, cytotoxic agents, nucleotides, antibodies, and non-protein micromolecules.

Preferably, the process of this invention involves a series of steps comprising:

- 10           a. sequentially coupling selected amino acids or amino acid analogs to a terminal amino acid or amino acid analog bound to a first resin support to form a first peptide segment-resin, the first peptide segment  
15           having about two to about one hundred amino acid residues;
- 20           b. covalently attaching a haloacyl moiety to the N-terminus of the first peptide segment-resin to form a haloacylpeptide segment bound to the first resin support;
- 25           c. cleaving the haloacylpeptide peptide segment from the first resin support;
- 30           d. sequentially coupling selected amino acids or amino acid analogs to a terminal amino acid or amino acid analog bound to a second resin support through a sulfur or selenium-containing bond to form a second peptide segment-resin, the second peptide segment having about two to about one hundred amino acid residues;

- 6 -

e. cleaving the second peptide segment-resin to form a second peptide segment having a thiol- or selenol-containing group at the C-terminus thereof; and

5

f. coupling the haloacylpeptide peptide segment and the second peptide segment to form a modified polypeptide.

10 The order of the sequence of steps a-b-c-d-e- is not critical to this invention. The sequence of steps a-b-c and the sequence of steps d-e may be conducted successively or separately. The entire sequence can be repeated in a chain-reaction manner.

15

Optionally, any reactive groups such as thiol that may be present in the peptide segments can be protected prior to step (f) and deprotected after step (f) is complete.

20

This invention, in its broadest sense, encompasses a biologically active protein comprising two molecular segments, each selected from peptides, pseudopeptides, or non-peptide linear molecules linked through a non-amido linkage to form a peptide or pseudopeptide backbone, wherein one of the segments contains at least one non-coded structural unit and the non-coded structural unit does not form a part of the non-amido linkage, provided that both segments are not a non-peptide linear molecule at the same time.


25

30

This invention further provides a modified protein represented by the formula:



- 7 -



wherein R and R' are the same or different and are each a residue of a peptide or pseudopeptide; and L represents a thiol ester or selenol ester linkage. Preferably, both R and R' comprise from about two to about one hundred amino acid residues.

The above objects and features of the invention will become more fully apparent from the description of the preferred embodiments in conjunction with the accompanying figures.

#### BRIEF DESCRIPTION OF THE FIGURES

15

FIGURE 1 shows a synthetic strategy for the total chemical synthesis of HIV-1 PR analogs in accordance with this invention where "A" schematically represents the coupling of the N-terminal segment HIV-1 PR (1-50, Gly<sup>51</sup> SH) and the C-terminal segment bromoacetyl (53-99) HIV-1 PR, and "B" schematically represents the coupling of the N-terminal segment HIV-1 PR (1-50, Cys<sup>51</sup> amide) and the same C-terminal segment.

25

FIGURE 2 shows an elution profile of aliquots taken at t = 0, 45 min., 3 h, and 48 h from the ligation mixtures containing HIV-1 PR (1-50, Gly<sup>51</sup>SH) and bromoacetyl (53-99) HIV-1 PR by the practice this invention using reverse phase HPLC (absorbance 214nm).

30

FIGURE 3A shows the ion spray mass spectrum of the purified ( $[NHCH_2COSCH_2CO]^{51-52}Aba^{67,95}$ ) HIV-1 PR where the labeled peaks represent a single molecular species differing in the number of excess protons.

- 8 -

FIGURE 3B shows the deconvoluted mass spectrum of the purified ( $[\text{NHCH}_2\text{COSCH}_2\text{CO}]^{51-52}$  Aba <sup>62,95</sup>) HIV-1 PR where the peak of the molecular weight of the enzyme is located at 10, 769 dalton.

5

FIGURE 4 shows an elution profile of aliquots taken at  $t = 0, 30 \text{ min},$  and  $120 \text{ min}$  from the ligation mixture containing HIV-1 PR(1-50, Cys amide) and bromoacetyl (53-99)] HIV-1 PR.

10

FIGURE 5 shows an elution profile of the hexapeptide Ac-Thr-Ile-Nle-Nle-Gln-Arg;  $\text{NH}_2$  before and after treatment with the ligation mixture taken at  $t = 3\text{h}$  as shown in FIGURE 2 using reverse phase HPLC with absorbance monitored at 214 nm, where the upper panel shows peaks before the treatment and the lower panel, peaks after the treatment, respectively.

15

FIGURE 6 shows a fluorogenic assay of aliquots of the ligation mixture taken at the times indicated as shown in FIGURE 2 where data points illustrating fluorescence units were read from continuous chart recorder tracings.

20

25

#### DESCRIPTION OF THE PREFERRED EMBODIMENTS

This invention is based on a conceptually novel approach to synthesizing large biomolecules such as proteins by a convergent type synthesis. Thus, when applied to the synthesis of modified proteins, this invention involves the coupling of at least two peptide segments to create a linkage which can be a non-amido bond. The peptide segments may be synthesized using solid phase peptide synthesis,

30

- 9 -

solution phase synthesis, or by other techniques known in the art including combinations of the foregoing methods.

5       The present process involves the steps of:

- 10           a.   providing a first segment having at least one non-coded structural unit, and a first chemoselective synthon at the terminal position thereof;
- 15           b.   providing a second segment optionally containing at least one non-coded structural unit, and a second chemoselective synthon at the terminal position thereof, the second chemoselective synthon being complementary to the first chemoselective synthon of the first segment; and
- 20           c.   ligating the first segment and the second segment, whereby the first synthon of the first segment and the second synthon of the second segment forms a non-peptide linkage, wherein the first segment and the second  
25           segment are each selected from peptides, pseudopeptides, or non-peptide linear molecules, provided that both segments are not non-peptide linear molecules at the same time.

30

Preferably, both of the first and second segments are peptides or pseudopeptides having from about two to about one hundred amino acid residues. More

- 10 -

preferably, the segments constitute from about forty to sixty amino acid residues.

5 Preferably, the non-peptide linkage formed is represented by one of the following linking moieties:

-CH<sub>2</sub>-S-, -CH<sub>2</sub>-Se-, -CO-S-, -CO-Se-, -CH<sub>2</sub>-NH-,  
or -CH(OH)-CH<sub>2</sub>-.

10 The present process allows at least two different non-coded structural units to be incorporated to the segments as well as allowing the same non-coded structural units to be incorporated at two sites.

15 In view of the broad and varied class of modified proteins which may be bound by this invention, all chemically modified proteins having such characteristics as defined above are deemed to be within the scope of this invention. However, for  
20 purposes of illustrative clarity and ease of comprehension, this invention will be described herein in more detail utilizing a modified HIV-1 protease as an embodiment. It should be understood that the use of this particular enzyme for descriptive purposes  
25 shall not restrict nor limit the use of other proteins or peptides.

As employed herein, the term "modified protein" is intended to include oligopeptides,  
30 oligopseudopeptides, polypeptides, pseudopolypeptides, and modified native proteins-synthetic or otherwise derived. The term "pseudopeptide" means a peptide where one or more peptide bonds are replaced by non-amido bonds such as ester or one or more amino acids

- 11 -

are replaced by amino acid analogs. The term "peptides" refers not only to those comprised of all natural amino acids, but also to those which contain unnatural amino acids or other non-coded structural units. The terms "peptides", when used alone, include pseudopeptides. The "modified proteins" have utility in many biomedical applications because of increased stability toward in vivo degradation, superior pharmacokinetics, and enhanced or diminished immunogenicity compared to their native counterparts.

HIV-1 protease (HIV-1 PR) is a virally-encoded enzyme which cuts polypeptide chains with high specificity and which is essential for the replication of active virions (N.E. Kohl, et al., *Proc. Natl. Acad. Sci., U.S.A.*, 85:4686, 1988). The 21,500 dalton HIV-PR molecule is made up of two identical 99 amino acid polypeptide chains.

Comparison of the crystal structures of the empty (A. Wlodawer, et al., *Science*, 245:616, 1989) and inhibitor-bound (for example, M. Miller, et al., *Science*, 246:1149, 1989) enzyme revealed that on binding a substrate-derived inhibitor the HIV-1 molecule undergoes significant conformational changes which are particularly pronounced in two exterior, functionally-important "flap" regions. From these crystallography studies it appears that peptide bonds in the flap regions of the HIV-1 PR polypeptide backbone are involved in the formation of  $\beta$ -sheet/ $\beta$ -turn structure, in the interaction which occurs between the two subunits of the active dimer at the tip of each flap in the enzyme-inhibitor (substrate), complex, and in hydrogen bonding interactions with

- 12 -

bound peptide inhibitors (and, presumably, substrates). Mutagenesis studies carried out with recombinant HIV-1 PR showed that the flap region is highly sensitive to changes in the amino acid sequence  
5 (D.D. Loeb, et al., *Nature*, 340:397, 1989).

These observations make the flap region especially interesting as a target for protein backbone modifications to investigate the role of peptide bond  
10 interactions in HIV-1 protease activity.

In the case of HIV-1 PR where modifications of the flap region are desired, this invention allows pseudo-peptide bonds to be introduced into the region. The  
15 Gly<sup>51</sup>-Gly<sup>52</sup> bond of HIV-1 PR is particularly preferred for bond manipulation for two reasons. First, glycine is the only achiral amino acid and therefore there is no concern about loss of optical purity; and second, the Gly<sup>51</sup>-Gly<sup>52</sup> bond is located near the middle of the  
20 99 amino acid HIV-1 PR monomer polypeptide chain. This means that the two peptide segments which are to be coupled are each about 50 residues in length.

The solid phase peptide synthesis method is generally  
25 described in the following references: Merrifield, J. Am. Chem. Soc., 88:2149, 1963; Barany and Merrifield, *In the Peptides*, E. Gross and J. Meinenhofer, Eds., Academic Press, New York, 3:285 (1980); S.B.H. Kent, *Annu. Rev. Biochem.*, 57:957 (1988). By the solid  
30 phase peptide synthesis method, a peptide of a desired length and sequence can be produced through the stepwise addition of amino acids to a growing peptide chain which is covalently bound to a solid resin

- 13 -

particle. Automated synthesis may be employed in this method.

Accordingly, these embodiments can be accomplished by the steps of:

1. sequentially coupling amino acids to a terminal amino acid bound to a resin support to form a peptide segment-resin; and
2. cleaving the peptide segment from the resin support.

In the preferred application of this method, the C-terminal end of the growing peptide chain is covalently bound to a  $-OCH_2$  PAM resin and amino acids having protected  $\alpha$ -amino groups are added in the stepwise manner indicated above. A preferred  $\alpha$ -amino protecting group is the tert-butyloxycarbonyl (BOC) group, which is stable to the condensation conditions and yet is readily removable without destruction of the peptide bonds or racemization of chiral centers in the peptide chain. At the end of the procedure the product peptide is cleaved from the resin, and any remaining protecting groups are removed by treatment under acidic conditions such as, for example, with a mixture of hydrobromic acid and trifluoroacetic acid, with trifluoromethane sulfonic acid or with liquified hydrogen fluoride.

In the case of HIV-1 protease, the C-terminal segment comprises the following amino acid sequence:  $F^{53}$  IKVRQYD<sup>60</sup> QIPVEICGHK<sup>70</sup> AIGTVLVGPT<sup>80</sup> PVNIIGRNLL<sup>90</sup> TQIGCTLNF<sup>99</sup>. The N-terminal

- 14 -

segment comprises the following amino acid sequence:

P<sup>1</sup>QITLWQRPL<sup>10</sup> VTIRIGGQLK<sup>20</sup> EALLDTGADD<sup>30</sup> TVLEEMNLPG<sup>42</sup>

KWKPKMIGGI<sup>50</sup>G<sup>51</sup>. The synthesis of these segments normally require about 15 hours at a standard cycle speed.

If desired, any amino acids in the above sequences can be replaced by amino acid analogs or amino acid mimetic compounds known in the art. Suitable amino acid substitutes include  $\beta$ -alanine, L- $\alpha$ -aminoisobutyric acid, L- $\alpha$ -amino-n-butyric acid (Aba), 3.4-dehydroproline, homoarginine, homocysteine, homoproline, homoserine, 3-mercaptopropionic acid, norleucine (Nle), penicillamine, pyroglutamic acid and sarcosine.

L-amino acids and D-amino acids may be used in this invention, particularly, D-amino acids are useful for the formation of a "reversible peptide sequence".

The "reversed" or "retro" peptide sequence as discussed above refers to that part of an overall sequence of covalently-bonded amino acid residues (or analogs or mimetics thereof) wherein the normal carboxyl-to-amino direction of peptide bond formation in the amino acid backbone has been reversed such that, reading in the conventional left-to-right direction, the amino portion of the peptide bond precedes (rather than follows) the carbonyl portion (see, generally, Goodman, M. and M. Chorev, *Accounts of Chem. Res.*, 12:423, 1979).

The "reversed" peptides are within the meaning of the "peptides" used throughout the specification. D-amino



- 15 -

acids, amino acid analogs and amino acid mimetic compounds are collectively referred to herein as "non-coded amino acids." Although the solid phase peptide synthesis theoretically enables one skilled in the art to prepare a peptide backbone of any length, the efficiency of coupling amino acids (addition of an amino acid in successive cycles) would necessarily limit the use of this technique when a peptide to be synthesized has greater than 150 residues. Preferably, for practical and economic reasons, the process of this invention employs from about two to about one hundred cycles.

Accordingly, the modified proteins obtained by the process of this invention have the general structure set forth in formula (1) and R and R' are as previously defined. If a larger peptide segment (containing greater than one hundred amino acid residues) is desired, such peptide segment may be available from naturally occurring proteins (native proteins) by enzymic or chemical degradation. Alternatively, the present process can be repeated with different ligation modes for building such large peptide segments.

An important feature of this invention is the bond formation linking two peptide segments (namely R and R'). A variety of bond forming reactions can be used to covalently link the two peptide segments. Representative combinations of such groups are carbonylthiol with halo to form a thiol ester bond between the two segments, carbonyl selenol with halo to form a seleno ester, or thiol with thiol to form a disulfide bond, thiol with halo to form a thioether

- 16 -

bond, selenol with halo to form a selenol ether, amino with isothiocyanate to form a thiourea bond, amino with aldehyde to form an imine bond which can be reduced to a carbon-nitrogen bond, thiol with maleimide to form a thioether bond, and gem-diol with boron to form oxygen-boron bonds, and hydroxyl with carboxyl to form an ester bond. Among these, preferred linkages are those already enumerated by way of their structure. A preferred linkage is a sulfur-containing or selenium-containing linkage which does not readily hydrolyze *in vivo* or which is more liable than an amido bond *in vivo*.

The most preferred linkage L is a thiol ester linkage. This linkage can be accomplished by first attaching a facile leaving group to a first peptide segment and by attaching carbonylthiol functionality to a second peptide segment followed by nucleophilic substitution where the sulfur nucleophile attacks the leaving group. Preferably, a haloacyl (e.g., haloacetyl) such as iodo, chloro, or bromoacetyl is attached to the N-terminus of the first peptide segment. The suitable haloacyl group may be straight or branched (substituted by alkyl). This step is conveniently carried out while the first peptide segment is still bound to a resin support.

To introduce a bromoacetyl group to the first peptide segment, suitable activated forms of bromoacetic acid may be employed in this invention. The particularly preferred agent for this purpose is bromoacetic anhydride.

- 17 -

- When the aforementioned C-terminal segment of HIV-1 PR is derivatized with the N-bromoacetylating agent, the unprotected N-terminus of the protected peptide segment on resin is condensed with bromoacetic anhydride to produce bromoacetyl (53-99) HIV-1 PR. Deprotection and release of the product peptide segment from the resin support can be accomplished by standard conditions (e.g., treatment with anhydrous HF containing 10% p-cresol at 0°C for several hours).
- 5      The product peptide segment is precipitated, dried by lyophilization, and purified, if desired, by reverse phase HPLC, according to standard techniques known in the art.
- 10     To introduce carbonylthiol functionality to the second peptide segment, its terminal carboxylic acid moiety may be converted to a carbonylthiol group.
- 15     When the aforementioned N-terminal segment of HIV-1 PR is to be derivatized accordingly, 4-[ $\alpha$ -(Boc-Gly-S) benzyl] phenoxyacetamidomethyl-resin is used as the resin support. If any amino acid or amino acid analog other than glycine is desired, that amino acid can be loaded on an aminomethyl-resin in the form of 4-[ $\alpha$ -(Boc-X-S) benzyl] phenoxyacetic acid wherein X represents the amino acid.
- 20     The N-terminal derivatized peptide segment can thus readily be prepared by the stepwise solid phase synthesis. The second product peptide segment is cleaved from the resin support, deprotected, and isolated in the manner described above.
- 25     The N-terminal derivatized peptide segment can thus readily be prepared by the stepwise solid phase synthesis. The second product peptide segment is cleaved from the resin support, deprotected, and isolated in the manner described above.
- 30     The N-terminal derivatized peptide segment can thus readily be prepared by the stepwise solid phase synthesis. The second product peptide segment is cleaved from the resin support, deprotected, and isolated in the manner described above.

- 18 -

The thiol ester linkage can be generated by coupling the two segments under normal ligation conditions. The formation of the thiol ester linkage is highly chemoselective and thus compatible with most reactive groups that may be present in the molecules.

A typical ligation reaction is carried out by mixing the unprotected N- and C- terminal segments in 6M guanidine hydrochloride buffer at about pH 3-6. In this buffer, the solubility of the unprotected peptide segments is very high, thus eliminating the major drawback of the prior art techniques having to use protected peptide fragments despite their limited solubility in ligation media. Other denaturants such as urea, detergents, and sodium dodecyl sulfate can be used as the ligation buffer.

The coupling of HIV-1 PR (1-50, Gly-SH) with bromoacetyl (53-99) HIV-1 PR is complete in several hours. The product peptide  $[(\text{NH CH}_2\text{COSCH}_2\text{CO})^{51-52}]$  HIV-1 PR can be isolated, purified and characterized by standard techniques.

When one of the peptide segments has any reactive groups which may interfere with the thiol ester formation, those groups may be protected by suitable protecting groups known in the art or the amino acids bearing such groups can be substituted by other amino acids or amino acid analogs which are reaction inert and yet do not adversely affect the biological activity of a product protein.

For example, in the Examples, the C-terminal segment of HIV-1 PR has two cysteine residues at positions 67

- 19 -

and 95. These cysteine positions have been shown to be replaceable by L- $\alpha$ -amino-n-butyric acid (Aba) without causing the loss of the enzymic activity of the native protease. In one embodiment of this invention, bromoacetyl [(53-99)Aba<sup>67,95</sup>] HIV-1 PR is used. Alternatively, to block a thiol group of the cysteine residues before the coupling reaction, the use of protecting groups that are compatible with ligation conditions are preferred. However, these precautions may not be necessary since, in present experience, the thiol ester group of the N-terminal segment attacks a bromoacetyl group even at low pH conditions where a thiol side chain of a cysteine residue is unreactive.

In another embodiment of the invention, the linkage L can be a thioether bond. This linkage can be accomplished in substantially the same manner as that described for the formation of the thiol ester linkage, except that thiol functionality is attached to a second peptide segment. This is conveniently accomplished by utilizing a thiol side chain of a cysteine residue. Thus, cysteine can be employed as the C-terminal amino acid of the second peptide segment (N-terminal segment). HIV-1 PR(1-50, Cys<sup>51</sup> amide) described herein represents one example of such derivatized second peptide segment where the cysteine's carboxylic acid is blocked as amide. When this peptide segment is subjected to the ligation conditions described earlier in the presence of bromoacetyl (53-99) HIV-1 PR, the coupling of the two peptide segments takes place rapidly to afford  $\{[NHCH(CONH_2)CH_2SCH_2CO]^{51-52}\}HIV-1$  PR. The ligated product protein can be isolated, purified, and

- 20 -

characterized by standard techniques. In a further embodiment of the invention, the linkage can be a selenium-containing bond such as selenol ester and seleno ether. In like manner, a selenocysteine can be used in place of cysteine.

The process of this invention is applicable to the synthesis of any biomolecules and is not limited to proteins insofar as two constituent fragments are available by chemical synthesis or other biosynthetic methods. In particular, the present process: (1) allows rapid synthesis of modified proteins; (2) avoids the use of protecting groups in at least the critical bond formation stage; and (3) incorporates into the protein backbone structural units such as D-amino acids and amino acid analogs.

This invention will be described in further detail below by way of the aforeindicated embodiments, but these embodiments should not be taken as limiting the scope of the invention.

- 21 -

EXAMPLE 1SYNTHESIS OF  $[\text{NHCH}_2\text{COSCH}_2\text{CO}]^{51-51}\text{Aba}^{67,95}$  HIV-1 PR

5 Two peptide segments, HIV-1 PR (1-50, Gly<sup>51</sup>SH)  
(PREPARATION 1) and bromoacetyl (53-99) HIV-1 PR  
(PREPARATION 2) were coupled by ligating the segments  
in 6M guanidine hydrochloride 0.1 M sodium phosphate  
at pH4.3. The segments were separately dissolved in  
10 ligation buffer at a concentration of 20 mg/ml.

The process of the ligation was followed by reverse  
phase HPLC on a Vydac C<sub>18</sub> column using a linear  
gradient of 30-60% buffer B (90% acetonitrile/0.09%  
15 trifluoroacetic acid) in buffer A (0.1%  
trifluoroacetic acid) in 30min. The flow rate was 1  
ml/min and absorbance was monitored at 214 nm.  
Results are shown in FIGURE 2.

20 The resulting ligated peptide was purified by reverse  
phase HPLC on a semipreparative Vydac C<sub>18</sub> column using  
various linear gradients of 90% acetonitrile/0.09% TFA  
in 0.1% aq TFA. The purity of the product peptide was  
checked by analytical HPLC as well as by ion spray  
25 mass spectrometry. The product peptide had the  
expected molecular weight of  $10.768.6 \pm 1.1$  Da (Calcd.  
for monoisotopic; 10.763.9 Da; (average) 10, 770.8  
Da). The mass spectra of the title peptide are shown  
in FIGURE 3A and 3B.

30

- 22 -

EXAMPLE 2SYNTHESIS OF [(NHCH(CO NH<sub>2</sub>) CH<sub>2</sub> 5CH<sub>2</sub> CO]<sup>51-52</sup>Aba<sup>67,95</sup>] HIV-1 PR

5 Two peptide segments HIV-1 PR (1-50, Cys<sup>51</sup> amide)  
(PREPARATION 1) and bromoacetyl (53-99) HIV-1 PR  
(PREPARATION 3) were coupled substantially according  
10 to the procedure of EXAMPLE 1, except that the pH was  
over 7.0. The ligation reaction was monitored by  
HCLP, with the results shown in FIGURE 3. After work-  
up and purification, the product peptide had the  
expected molecular weight of 10,800.31 ± 0.75 Da  
(Calcd. for monoisotopic 10,792.9 Da; (average)  
15 10,799.8 Da).

EXAMPLE 3ENZYMATIC ACTIVITY OF [(NHCH<sub>2</sub>COSCH<sub>2</sub>CO)<sup>51-52</sup>Aba<sup>67,95</sup>] HIV-1 PR

20 An aliquot of the ligation reaction mixture after 3  
hours ligation (EXAMPLE 1) was treated with a peptide  
substrate (1 mg/ml) having the sequence of Ac-Thr-Ile-  
25 Nle-Nle-Gln-Arg. NH<sub>2</sub> (Nle:L-norleucine) at pH 6.5.  
Reaction was monitored by reverse phase HPLC using a  
Vydac C18 column with the results shown in FIGURE 5.  
In FIGURE 5 the upper panel shows the peptide  
substrate peaks before treatment. The lower panel  
30 shows the peaks of cleavage products after 15 min.  
treatment.

The cleavage products were separated by reverse phase